

Path regularity of Gaussian processes via small deviations

Frank Aurzada

May 20, 2009

Abstract

We study the a.s. sample path regularity of Gaussian processes. To this end we relate the path regularity directly to the theory of small deviations. In particular, we show that if the process is n -times differentiable then the exponential rate of decay of its small deviations is at most $\varepsilon^{-1/n}$. We also show a similar result if n is not an integer.

Keywords: Small deviation; small ball probability; Gaussian process; sample path regularity

2000 Mathematics Subject Classification: 60G15; 60F99

Running Head: Path regularity via small deviations

1 Introduction

The small deviation problem, also called small ball problem, consists in determining the rate of increase of the quantity

$$-\log \mathbb{P} [\|X\| \leq \varepsilon], \quad \text{as } \varepsilon \rightarrow 0. \quad (1)$$

Here, X is a random variable with values in a normed space $(E, \|\cdot\|)$.

This problem has several connections to approximation quantities for stochastic processes. We refer to [LS01] for an overview of the field and links to applications and to [Lif08] for a regularly updated list of references. Recently, several articles have focused on the small deviation problem for integrated Gaussian processes, [KS98, CL03, GHT03, GHLT03, FT04, NN04a, LS05, Gao08]. This is mainly due to the connection of the problem to the spectral asymptotics of certain boundary value problems.

In some sense, this paper also considers integrated processes. However, we do not aim at finding the rate in (1) for Gaussian processes, but rather at showing that this rate is directly related to the sample path regularity of the process.

The idea that the small deviation rate encodes the smoothness properties of the Gaussian process has been present in many articles on small deviations. However, it seems that no concrete results are available that relate the small deviation rate *directly* to smoothness properties of the process, e.g. differentiability. The aim of this article is to provide this direct link.

In particular, we are going to show (Corollary 9 below) that if a Gaussian process is n -times differentiable then for its small deviation rate

$$-\log \mathbb{P} \left[\|X\|_{L_\infty[0,1]} \leq \varepsilon \right] \leq c \varepsilon^{-1/n}, \quad \text{as } \varepsilon \rightarrow 0.$$

We also show a similar result if n is not an integer. As we shall see, this provides sharp criteria for the path regularity of Gaussian processes. Note that for example for Brownian motion the small deviation rate is ε^{-2} , which corresponds to being Hölder continuous up to $\frac{1}{2}$. A consequence is that for a $\mathcal{C}^\infty$ process we have that, for any $\delta > 0$,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^\delta \left(-\log \mathbb{P} \left[\|X\|_{L_\infty[0,1]} \leq \varepsilon \right] \right) = 0.$$

These results, combined with Li's weak decorrelation inequality [Li99], have one further consequence (Corollary 21 below): If the Gaussian process X can be represented as $X = Y + Z$, with Y and Z not necessarily independent, and Z is smooth enough then X and Y have the same small deviation order. This can be used to show that the small deviation order of smoother 'remainder terms' does not matter, as will be demonstrated with some examples.

For showing the relation between path regularity and small deviations we employ a result developed by Chen and Li in [CL03]. They show the following (Theorem 1.2 in [CL03]).

Proposition 1. *Let X be a centered Gaussian random variable with values in some separable Banach space $(E, \|\cdot\|)$. Let $\mathcal{H}$ be the reproducing kernel Hilbert space of X and denote by $|\cdot|_{\mathcal{H}}$ the norm induced by the inner product in $\mathcal{H}$. Let Y be another Gaussian random variable in $(E, \|\cdot\|)$, not necessarily independent of X . Then, for any $\lambda, \varepsilon > 0$,*

$$\mathbb{P} [\|Y\| \leq \varepsilon] \geq \mathbb{P} [\|X\| \leq \lambda\varepsilon] \mathbb{E} \exp \left(-\frac{\lambda^2}{2} |Y|_{\mathcal{H}}^2 \right). \quad (2)$$

In particular, let E be a space of functions from $[0, 1]$ to $\mathbb{R}$ and let B be Brownian motion. Then

$$\mathbb{P} [\|Y\| \leq \varepsilon] \geq \mathbb{P} [\|B\| \leq \lambda\varepsilon] \mathbb{E} \exp \left(-\frac{\lambda^2}{2} \|Y'\|_{L_2[0,1]}^2 \right), \quad (3)$$

for any $\lambda, \varepsilon > 0$ and any Gaussian random variable Y with values in E , where Y' is the derivative of Y .

This result was used to derive the small deviation rate for the m -fold integrated Brownian motion in [CL03]. The procedure is as follows: Let Y be integrated Brownian motion. Knowing the small deviation rate of Brownian motion Y' gives the rate of the Laplace transform on the right-hand side in (3), when $\lambda \rightarrow \infty$. The small deviation probability of Brownian motion B w.r.t. $\|\cdot\|$ on the right hand side in (3) is known as well. This gives a lower bound for the small deviation probability of integrated Brownian motion Y for basically any norm $\|\cdot\|$. The procedure can be iterated. On the other hand, the upper bound for the small deviation rate of Y e.g. for the L_∞ -norm can be obtained simply by comparison to the easier L_2 -norm.

The focus of the present note is

- to formulate the idea from [CL03] in a general framework (Section 2.1), and to extend it to fractional derivatives (Section 2.2),
- to show that this leads to information on the a.s. path regularity of the Gaussian process under consideration (Section 3),
- to study a conditional version of (2) that can be applied in particular to stable processes (Section 4.1), and
- to investigate relations to other questions (Sections 4.2, 4.3, 4.4) and concrete examples (Section 4.5).

In this paper, we let X be a real-valued, centered Gaussian process indexed by $[0, 1]$ with $X(0) = 0$ a.s. The restriction $X(0) = 0$ is for simplicity only. We use $\sim$, $\gtrsim$, and $\lesssim$ for strong asymptotics, i.e. $f \gtrsim g$ or $g \lesssim f$ if $\limsup f/g \leq 1$, $f \sim g$ if $\lim f/g = 1$, whereas $\approx$, $\gtrsim$, and $\lesssim$ stand for weak asymptotics, i.e. $f \gtrsim g$ or $g \lesssim f$ if $\limsup f/g < \infty$ and $f \approx g$ if $f \gtrsim g$ and $f \lesssim g$. We frequently use $1/0 = \infty$ and $1/\infty = 0$.

2 Results for small deviations

2.1 First results

Our first theorem concretizes the method used in [CL03]. Here, X' denotes the derivative of X .

Theorem 2. *Let $0 < \tau \leq \infty$, $\theta \in \mathbb{R}$, and $1 \leq p \leq \infty$. Then*

$$-\log \mathbb{P} \left[\|X'\|_{L_2[0,1]} \leq \varepsilon \right] \lesssim \kappa \varepsilon^{-1/\tau} |\log \varepsilon|^\theta \quad (4)$$

implies

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \lesssim C \varepsilon^{-1/(\tau+1)} |\log \varepsilon|^{\theta\tau/(\tau+1)},$$

where $C = C(\kappa, \kappa_p)$ and κ_p is the small deviation constant of Brownian motion w.r.t. the L_p -norm.

The proof goes along the lines outlined above; we skip it since it is included in the more general Theorem 6 below.

Remark 3. We remark that the constant $C = C(\kappa, \kappa_p)$ can be computed explicitly. The constant is, in general, not the correct small deviation constant for X w.r.t. L_p -norm. For $\tau = \infty$ (explicitly permitted in the above theorem) we get $C = \kappa$. The case $\tau = 0$ is of special interest; and we treat it in Corollary 9.

Using the same idea as in Theorem 2 we can obtain a converse estimate.

Theorem 4. Let $0 \leq \gamma < 1$, $\delta \in \mathbb{R}$ and $1 \leq p \leq \infty$. Then

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \gtrsim C \varepsilon^{-\gamma} |\log \varepsilon|^\delta,$$

implies

$$-\log \mathbb{P} \left[\|X'\|_{L_2[0,1]} \leq \varepsilon \right] \gtrsim \kappa \varepsilon^{-\gamma/(1-\gamma)} |\log \varepsilon|^{\delta/(1-\gamma)}$$

where $\kappa = \kappa(C, \kappa_p)$ and κ_p is the small deviation constant of Brownian motion w.r.t. the L_p -norm.

The cases $\gamma = 0$ (included above) and $\gamma = 1$ (treated in Corollary 9) are of special interest.

2.2 Fractional derivatives

After demonstrating the method from [CL03], we now extend the idea to a more subtle situation. Here we define the fractional derivative as follows (cf. [SKM93]). Recall that we work with processes with $X(0) = 0$. For a given function F with $F(0) = 0$ and $M > 0$, we set

$$\frac{d^M F}{dt^M}(t) = f(t) \quad :\Leftrightarrow \quad F(t) = \int_0^t (t-s)^{M-1} f(s) ds. \quad (5)$$

We stress that the M -th derivative of F (M integer) coincides with $d^M F/dt^M$. If there is no ambiguity we also write for simplicity

$$F^{(M)} = \frac{d^M F}{dt^M}.$$

Now we get an analog to Theorem 2.

Theorem 5. *Let $0 < \tau \leq \infty$, $\theta \in \mathbb{R}$, $1 \leq p \leq \infty$, and $M > 0$. Then*

$$-\log \mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]} \leq \varepsilon \right] \lesssim \kappa \varepsilon^{-1/\tau} |\log \varepsilon|^\theta$$

implies

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \lesssim C \varepsilon^{-1/(\tau+M)} |\log \varepsilon|^{\theta\tau/(\tau+M)},$$

where $C = C(\kappa, \kappa_p^M)$ and κ_p^M is the small deviation constant of a standard Riemann-Liouville process R^H (cf. (7) below) with $H = M - 1/2$ w.r.t. the L_p -norm.

The proof goes along the same lines as the one of Theorem 2. The only difference is that we use the Riemann-Liouville process R^H instead of Brownian motion B . An analog to Theorem 4 also holds true.

For $\tau = \infty$ (explicitly permitted in the above theorem) we get $C = \kappa$. Remark 3 applies accordingly.

2.3 General result for translation invariant, self-similar, pseudo-additive norms

In this section, we give the most general result for small deviations. For this purpose, we recall from [LS05] the notion of $\|\cdot\|$ being a translation invariant, β -self-similar, and p -pseudo additive functional semi-norm, for short $\|\cdot\| \in \mathbf{N}(\beta, p)$. Here, we require that $\|\cdot\|$ be a true norm of a separable Banach space. Instead of rewriting the definition from [LS05] we recall that the notion includes, for example, $L_p[0, 1]$ -norms ($\in \mathbf{N}(-1/p, p)$), η -Hölder norms ($\in \mathbf{N}(\eta, \infty)$), and the p -variation norm ($\in \mathbf{N}(0, p)$). Possibly it also includes certain Besov and Sobolev norms, see remarks in [LS05]. Since $X(0) = 0$, we are sure that we deal with a norm rather than a semi-norm.

Theorem 6. *Let $\|\cdot\| \in \mathbf{N}(\beta, p)$, $M > 0$, $0 < \tau \leq \infty$, and $\theta \in \mathbb{R}$. Then*

$$-\log \mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]} \leq \varepsilon \right] \lesssim \kappa \varepsilon^{-1/\tau} |\log \varepsilon|^\theta \tag{6}$$

implies

$$-\log \mathbb{P} [\|X\| \leq \varepsilon] \lesssim C \varepsilon^{-1/(\tau+M-\beta-1/p)} |\log \varepsilon|^{\theta\tau/(\tau+M-\beta-1/p)},$$

where $C = C(\kappa, \kappa_{\|\cdot\|}^M)$ and $\kappa_{\|\cdot\|}^M$ is the small deviation constant of a standard Riemann-Liouville process R^H (cf. (7) below) with $H = M - 1/2$ w.r.t. $\|\cdot\|$.

Proof: Note that (6) implies

$$-\log \mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]}^2 \leq \varepsilon \right] \lesssim \frac{\kappa}{2^\theta} \varepsilon^{-1/(2\tau)} |\log \varepsilon|^\theta.$$

By de Bruijn's Tauberian theorem (Theorem 4.12.9 in [BGT89]), this implies

$$-\log \mathbb{E} \exp \left(-\frac{\lambda^2}{2} \|X^{(M)}\|_{L_2[0,1]}^2 \right) \lesssim K \lambda^{1/(\tau+1/2)} |\log \lambda|^{\theta\tau/(\tau+1/2)},$$

when $\lambda \rightarrow \infty$. Here K can be computed explicitly from κ . We only remark that $\kappa = K$ for $\tau = \infty$. Note that the set

$$\mathcal{H} := \left\{ g : [0, 1] \rightarrow \mathbb{R} \mid g(t) = \int_0^t (t-s)^{H-1/2} f(s) ds, f \in L_2[0, 1], f(0) = 0 \right\}$$

with $H = M - 1/2$ and norm

$$\|g\|_{\mathcal{H}} = \|f\|_{L_2[0,1]} = \|g^{(M)}\|_{L_2[0,1]}$$

is the reproducing kernel Hilbert space ([Lif95]) of the Riemann-Liouville process

$$R^H(t) := \int_0^t (t-s)^{H-1/2} dB(s), \quad (7)$$

where B is a Brownian motion. Therefore, Proposition 1 implies that

$$\mathbb{P} [\|X\| \leq \varepsilon] \geq \mathbb{P} [\|R^H\| \leq \lambda\varepsilon] \mathbb{E} \exp \left(-\frac{\lambda^2}{2} \|X^{(M)}\|_{L_2[0,1]}^2 \right). \quad (8)$$

We can use the results for the Riemann-Liouville process from [LS05], which yield that

$$-\log \mathbb{P} [\|R^H\| \leq \lambda\varepsilon] \sim \kappa_{\|\cdot\|}^M (\lambda\varepsilon)^{-1/(H-\beta-1/p)}, \quad (9)$$

as long as $\lambda\varepsilon \rightarrow 0$. Set $\gamma := 1/(H - \beta - 1/p)$. We use this with

$$\lambda := D\varepsilon^{-(\tau+1/2)/(1/\gamma+\tau+1/2)} |\log \varepsilon|^{-\tau\theta/(\gamma(1/\gamma+\tau+1/2))},$$

where D is some constant. This gives

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon^{1/(1/\gamma+\tau+1/2)} |\log \varepsilon|^{\tau\theta/(1/\gamma+\tau+1/2)} \left(-\log \mathbb{P} [\|X\|_{L_p[0,1]} \leq \varepsilon] \right) \\ \leq \inf_{D>0} (\kappa_{\|\cdot\|}^M D^{-\gamma} + K D^{1/(\tau+1/2)}) =: C = C(\kappa, \kappa_{\|\cdot\|}^M). \end{aligned}$$

□

Analogously to Theorem 4 we can prove the following in the general setup.

Theorem 7. Let $0 \leq \gamma < 1$, $\delta \in \mathbb{R}$, and $\|\cdot\| \in \mathbf{N}(\beta, p)$. Then

$$-\log \mathbb{P} [\|X\| \leq \varepsilon] \gtrsim C \varepsilon^{-\gamma} |\log \varepsilon|^\delta,$$

implies

$$-\log \mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]} \leq \varepsilon \right] \gtrsim \kappa \varepsilon^{-1/(1/\gamma - M + \beta + 1/p)} |\log \varepsilon|^{\delta/(\gamma(1/\gamma - M + \beta + 1/p))}.$$

The proof is analogous to the one of Theorem 6.

3 Results for the path regularity

We now come to the mentioned results on the path regularity. In many articles on small deviations for Gaussian processes the authors mention that one can read off the path regularity from the small deviation results. However, to the knowledge of the author, no concrete result has been available so far.

The results on the path regularity follow from a modification of the proof of Theorem 6. Corresponding to $\tau = 0$ in Theorem 6 one obtains the following.

Theorem 8. Let $X^{(M)} \in L_2[0, 1]$ a.s. with $M > 0$. Then

(i) for any $1 \leq p \leq \infty$,

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \asymp \varepsilon^{-1/M}.$$

(ii) Furthermore, if $\|\cdot\| \in \mathbf{N}(\beta, p)$ then

$$-\log \mathbb{P} [\|X\| \leq \varepsilon] \asymp \varepsilon^{-1/(M - \beta - 1/p)}.$$

Proof: We only have to show (ii). The case of L_p -norms, part (i), is only a special case.

If $X^{(M)} \in L_2[0, 1]$ a.s. then there is a $K > 0$ such that

$$\mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]} \leq K \right] =: q > 0.$$

Using the Markov inequality, we obtain

$$q = \mathbb{P} \left[\|X^{(M)}\|_{L_2[0,1]} \leq K \right] = \mathbb{P} \left[e^{-\frac{\lambda^2}{2}} \|X^{(M)}\|_{L_2[0,1]}^2 \geq e^{-\frac{\lambda^2 K^2}{2}} \right] \leq \frac{\mathbb{E} \exp(-\frac{\lambda^2}{2} \|X^{(M)}\|_{L_2[0,1]}^2)}{\exp(-\frac{\lambda^2 K^2}{2})}.$$

Concerning the reproducing kernel Hilbert space with argue as in (8). Therefore, Proposition 1 implies that, setting $H := M - 1/2$ and $1/\gamma := H - \beta - 1/p$, we have

$$\mathbb{P} [\|X\| \leq \varepsilon] \geq \mathbb{P} [\|R^H\| \leq \lambda\varepsilon] \mathbb{E} \exp \left(-\frac{\lambda^2}{2} \|X^{(M)}\|_{L_2[0,1]}^2 \right) \geq e^{-c(\lambda\varepsilon)^{-\gamma}} q e^{-\frac{\lambda^2 K^2}{2}},$$

where we used the result for Riemann-Liouville processes (9) for the first term. Setting $\lambda := \varepsilon^{-\gamma/(2+\gamma)}$, this gives

$$-\log \mathbb{P} [\|X\| \leq \varepsilon] \lesssim \varepsilon^{-2\gamma/(2+\gamma)} = \varepsilon^{-1/(M-\beta-1/p)}.$$

□

In particular, in the case of integer derivatives, we obtain the following.

Corollary 9. *If the process X is n -times differentiable with $X^{(n)} \in L_2[0, 1]$ then*

$$-\log \mathbb{P} [\|X\|_{L_p[0,1]} \leq \varepsilon] \lesssim \varepsilon^{-1/n}.$$

If the process is C^∞ then

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^\delta \left(-\log \mathbb{P} [\|X\|_{L_p[0,1]} \leq \varepsilon] \right) = 0, \quad (10)$$

for all $\delta > 0$.

A close look at Theorem 8 and Corollary 9 reveals the following interesting interpretation: if the process has a certain path regularity – $X^{(M)}$ is in L_2 – then the small deviation probability cannot be too small – the logarithmic small deviation probability cannot grow faster than $\varepsilon^{-1/M}$. Conversely, if the small deviations grow too fast, then the path of the process cannot be too regular.

This is the first very concrete result of the intuitive fact that path regularity and small deviations for Gaussian processes are closely connected. We stress that beyond $\mathcal{C}^\infty$ the small deviation asymptotics has still distinct rates giving additional information.

Remark 10. We remark that Theorem 8 and Corollary 9 usually do *not* give the precise small deviation order even though we know the precise path regularity. Typically, $X^{(M)} \in L_2$ for all $M > M_0$. And thus the theorem yields

$$-\log \mathbb{P} [\|X\|_{L_p[0,1]} \leq \varepsilon] = \varepsilon^{-1/M_0 - o(1)}.$$

However, in many cases one finds that the above holds without the $o(1)$ term:

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \approx \varepsilon^{-1/M_0}.$$

One can think of Brownian motion itself (where $M_0 = 1/2$) or Riemann-Liouville processes. So, one can only hope to determine the small deviation asymptotics from the path regularity up to an $o(1)$ term.

4 Related questions

4.1 Remarks on stable processes

One may ask whether it is possible to extend the above results beyond the setup of Gaussian processes. This is indeed possible.

Since symmetric α -stable processes (in the sense of [ST94]) can be represented as conditionally Gaussian processes, the main tool used in the proofs, Theorem 1.2 in [CL03], can be transferred. Therefore, Theorems 2, 4, 5, 6, and 7 as well as Theorem 8 and Corollary 9 hold true also for symmetric α -stable processes.

However, it is easy to construct stable processes such that Theorem 8 and Corollary 9 do not give sharp results. This can be seen from the following example.

Example 11. Let X be a subfractional Brownian motion, i.e.

$$X(t) = A^{1/2} B^H(t), \quad t \geq 0,$$

where A is a strictly positive $\alpha/2$ -stable random variable and B^H is a fractional Brownian motion independent of A . Then obviously X has exactly the same a.s. path properties as B^H . So, somehow one might want to expect that X should have the same small deviation order as B^H . This is not the case, as shown by Samorodnitsky [Sam98]:

$$-\log \mathbb{P} \left[\|X\|_{L_\infty[0,1]} \leq \varepsilon \right] \approx \varepsilon^{-\frac{1}{1/\alpha - 1/2 + H}} \quad \text{vs.} \quad -\log \mathbb{P} \left[\|B^H\|_{L_\infty[0,1]} \leq \varepsilon \right] \sim C \varepsilon^{-1/H}.$$

This underlines that, partially, the small deviation rate is due to the fluctuations of the process, i.e. the path regularity, partially it is due to the (in this case: heavy) tail behaviour of the process, i.e. the amplitudes of the fluctuations.

4.2 Eigenvalues of the covariance operator

We recall that small deviation probabilities in L_2 -norm are closely connected (see e.g. [Naz03, NN04a, NN04b] and references therein) to the eigenvalues of the integral equation:

$$\lambda_n f_n(t) = \int_0^1 R(t, s) f_n(s) ds, \quad n \geq 1,$$

where $R(t, s) = \mathbb{E}X(t)X(s)$ is the covariance kernel of the Gaussian process X . Our results imply the following.

Corollary 12. *Let $\tau > 0$ and $\theta \in \mathbb{R}$. Let (λ_n) , (λ_n^1) be the sequence of eigenvalues of the operators given by the kernels*

$$R(t, s) \quad \text{and, respectively,} \quad \frac{\partial^2 R}{\partial t \partial s}(t, s)$$

Then

$$\lambda_n^1 \lesssim n^{-\tau} (\log n)^\theta$$

implies

$$\lambda_n \lesssim n^{-\tau-2} (\log n)^\theta.$$

A similar corollary can be obtained for fractional derivatives.

4.3 Quantization

We recall from [DFMS03, GLP03] that small deviations for Gaussian processes are closely related to the quantization problem. Let $D(r|X, \|\cdot\|)$ be the quantization error of the process X w.r.t. the distortion given by the norm $\|\cdot\|$, i.e. for a normed space $(E, \|\cdot\|)$, $s > 0$, and $r > 0$,

$$D(r|X, \|\cdot\|) := \inf \left\{ \left(\mathbb{E} \min_{a \in \mathcal{C}} \|X - a\|^s \right)^{1/s} : \mathcal{C} \subseteq E, \log \#\mathcal{C} \leq r \right\}.$$

The idea behind this quantity is that a random signal $X = X(\omega)$ has to be encoded; as a code one can use a minimizer $a(\omega) \in \mathcal{C}$ (minimizing $\min_{a \in \mathcal{C}} \|X(\omega) - a\|$); and if $\mathcal{C}$ was chosen close to optimal, this procedure gives a lowest possible mean error of the coding.

From the above results on small deviations and the connection established in [DFMS03, GLP03] one can obtain the following corollaries. The first gives the flavour of the more general result.

Corollary 13. *Let X be a centered Gaussian process on $[0, 1]$ that is differentiable and such that $X' \in L_2[0, 1]$ a.s. Let $\tau > 0$ and $\theta \in \mathbb{R}$. Then*

$$D(r|X', \|\cdot\|_{L_2[0,1]}) \lesssim r^{-\tau}(\log r)^\theta$$

implies

$$D(r|X, \|\cdot\|_{L_p[0,1]}) \lesssim r^{-(\tau+1)}(\log r)^\theta$$

for any $1 \leq p \leq \infty$.

This result can be used in the following way: In order to find the quantization rate of X w.r.t. the L_p -norm, it suffices to find a good estimate for the (easier) quantization problem for X' w.r.t. the L_2 -norm. Unfortunately, this connection is *not constructive*, so it is not clear how to obtain a good quantizer for X in L_p given that one has a good quantizer for the derivative X' w.r.t. L_2 distortion. It would be interesting to find a constructive proof of this fact.

Now we come to a more general corollary for the quantization error.

Corollary 14. *Let X be a Gaussian process on $[0, 1]$, $\|\cdot\| \in \mathbf{N}(\beta, p)$, $M > 0$, $\tau > 0$, and $\theta \in \mathbb{R}$. Then*

$$D(r|X^{(M)}, \|\cdot\|_2) \lesssim r^{-\tau}(\log r)^\theta$$

implies

$$D(r|X, \|\cdot\|) \lesssim r^{-(\tau+M-\beta-1/p)}(\log r)^\theta.$$

In particular, our results have the following corollary corresponding to the case $\tau = 0$ above.

Corollary 15. *Let X be a Gaussian process on $[0, 1]$, $\|\cdot\| \in \mathbf{N}(\beta, p)$, and $M > 0$. If $X^{(M)} \in L_2[0, 1]$ a.s. then*

$$D(r|X, \|\cdot\|) \lesssim r^{-(M-\theta-1/p)}.$$

Results that are very similar to Corollary 15 for *not necessarily Gaussian processes* were obtained in Lemma 2.1 in [Der08] (also see [DV04]). The technique used there is based on entropy numbers of embeddings and is thus more robust than the Gaussian techniques employed here. However, it seems only possible to use these techniques in the sense of ‘remainder terms’ as in Corollary 15, not the way we use it in Theorem 6.

Furthermore, we refer to [LP08] for similar results translating the mean path regularity (given by the behaviour of the covariance function in the Gaussian case) into estimates for the quantization error. There, wavelet representations for the process are used to obtain good quantizers. It seems possible that this approach can help to extend the results of this paper from the Gaussian setup to other processes.

Remark 10 applies accordingly to Corollary 15. However, we stress that Corollary 14 is sharp.

4.4 Entropy numbers of operators

Via the connections between small deviations and entropy numbers for linear operators ([KL93], [LL99], [AILvZ08]) we can obtain the following corollaries for operators related to Gaussian processes.

Let $M > 0$ and let $K : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be a measurable function with

$$K(t, \cdot) \in L_2[0, 1] \quad \text{and} \quad \frac{\partial^M K}{\partial t^M}(t, \cdot) \in L_2[0, 1],$$

where the fractional derivative is as defined in (5).

We consider the following operators

$$\begin{aligned} u : L_2[0, 1] &\rightarrow L_p[0, 1], & (uf)(t) &:= \int_0^1 K(t, s)f(s) ds \\ u_M : L_2[0, 1] &\rightarrow L_2[0, 1], & (u_M f)(t) &:= \int_0^1 \frac{\partial^M K}{\partial t^M}(t, s)f(s) ds. \end{aligned}$$

For a linear operator $u : E \rightarrow F$ between Banach spaces E and F one defines the entropy numbers as follows [CS90]:

$$e_n(u : E \rightarrow F) = \inf\{\varepsilon > 0 : \exists f_1, \dots, f_{2^{n-1}} \in F \forall x \in E, \|x\| \leq 1 : \|u(x) - f\| \leq \varepsilon\}.$$

The following theorem relates the (easy) $L_2[0, 1] \rightarrow L_2[0, 1]$ entropy numbers of u_M to the (more difficult) $L_2[0, 1] \rightarrow L_p[0, 1]$ entropy numbers of u .

Theorem 16. *Let $\tau > 0$, $\theta \in \mathbb{R}$, and $1 \leq p \leq \infty$. Set $E = L_p[0, 1]$. If*

$$e_n(u_M : L_2[0, 1] \rightarrow L_2[0, 1]) \lesssim n^{-1/2-\tau}(\log n)^\theta$$

then

$$e_n(u : L_2[0, 1] \rightarrow L_p[0, 1]) \lesssim n^{-1/2-\tau-M}(\log n)^\theta.$$

This theorem has the following interpretation: the regularity of the kernel (namely, the fact that $\partial^M K / \partial t^M$ is in $L_2[0, 1]$) can be used to translate an (easy) $L_2 \rightarrow L_2$ estimate for u_M into an $L_2 \rightarrow L_p$ estimate for u .

We remark that this theorem is *not* an embedding type theorem, since it holds for all $p \geq 2$ and $1 \leq p \leq 2$.

Remark 17. Let us mention that one can treat the cases $\tau = 0$ and $\tau = \infty$ accordingly.

First we look at the case $\tau = 0$. If for some $\theta > 1/2$

$$e_n(u_M : L_2[0, 1] \rightarrow L_2[0, 1]) \lesssim n^{-1/2}(\log n)^{-\theta}$$

then

$$e_n(u : L_2[0, 1] \rightarrow L_p[0, 1]) \lesssim n^{-1/2-M}.$$

The case $\tau = \infty$ can also be treated as above. Namely, for $\gamma > 0$

$$|\log e_n(u_M : L_2[0, 1] \rightarrow L_2[0, 1])| \gtrsim n^{-\gamma} \quad \Rightarrow \quad |\log e_n(u : L_2[0, 1] \rightarrow L_p[0, 1])| \gtrsim n^{-\gamma}.$$

Similar estimates can be obtained for other norms. In particular, if the $L_p[0, 1]$ -norm is replaced by a norm $\|\cdot\| \in \mathbf{N}(\beta, p)$ then M in the assertions has to be replaced by $M - \beta - 1/p$.

4.5 Examples

The first example is the integrated fractional Brownian motion.

Example 18. Consider the case when X is an integrated fractional Brownian motion. Then one obtains

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \lesssim \varepsilon^{-\frac{1}{H+1}}.$$

See e.g. [NN04a]. The lower bound can be obtained by comparing $\|X\|_{L_p[0,1]}$ to the L_2 -norm of X itself, which has the same order. Note that the H comes from fractional Brownian motion (and there from being Hölder up to H) and the 1 from one integration.

Consider the case where X is m -times integrated fractional Brownian motion. Then $X^{(m)}$ is precisely fractional Brownian motion and Theorem 5 states that a lower bound for the L_2 small deviations of fractional Brownian motion can be translated into a lower bound for the L_p small deviations of m -times integrated fractional Brownian motion. The result is

$$-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \lesssim \varepsilon^{-1/(H+m)},$$

which is the correct order, cf. [NN04a] and Theorem 1.3 in [CL03] for the Brownian case. The lower bound follows by comparison to the L_2 small deviations of X itself. This method was already used in [CL03] for the Brownian case $H = 1/2$.

The second example concerns a conjecture by Lifshits and Simon [LS05].

Example 19. Set

$$R^H(t) := \int_0^t (t-s)^{H-1/2} dB(s), \quad B^H(t) := R^H + \int_{-\infty}^0 (t-s)^{H-1/2} - (-s)^{H-1/2} dB(s), \quad (11)$$

where B is a Brownian motion. Then B^H is a fractional Brownian motion and R^H is a Riemann-Liouville process. Then one can look at the difference process $M = R^H - B^H$. This process is $\mathcal{C}^\infty$; and thus we have by Corollary 9

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^\delta \left(-\log \mathbb{P} \left[\|X\|_{L_p[0,1]} \leq \varepsilon \right] \right) = 0, \quad \text{for any } \delta > 0,$$

as conjectured by Lifshits and Simon [LS05]. In particular, this makes the proof of Theorem 12 in [LS05] redundant. The same result is true if B^H is a so-called linear fractional stable motion and R^H is a symmetric stable Riemann-Liouville process, see [LS05].

The next example is of similar type.

Example 20. We consider so-called stable convolutions [KMR06], i.e. processes

$$X(t) = \int_0^t f(t-s) dB(s), \quad t \geq 0,$$

where B is Brownian motion and f is a smooth function except possibly at zero, where we assume $f(x) = x^{H-1/2}g(x)$ with $H > 0$ and a function g with $g(0) = 1$. This setup was studied in [AS07]. The goal is to show that X has the same small deviation order (and path regularity) as a Riemann-Liouville process R^H .

We assume that g has the Taylor expansion $g(x) = \sum_{n=0}^\infty a_n x^n$, $0 \leq x \leq 1$. Let R^H be a Riemann-Liouville process as defined in (11). Then we can represent X as

$$\begin{aligned} X(t) &= \int_0^t (t-s)^{H-1/2} g(t-s) dB(s) \\ &= \int_0^t (t-s)^{H-1/2} dB(s) + \sum_{n=1}^\infty \int_0^t (t-s)^{H-1/2+n} a_n dB(s) = R^H(t) + \sum_{n=1}^\infty a_n R^{H+n}(t). \end{aligned}$$

Here the processes R^{H+n} , $n \geq 0$, are *not* independent but rather obtained by integrating the same Brownian motion as in (11).

Now the weak decorrelation inequality of Li [Li99] implies the following. In order to show that X and R^H have the same small deviation rate it suffices to know that the difference process $X - R^H$ has a lower small deviation order than R^H . However, this follows easily from Theorem 8, since we know that $X - R^H$ is smoother than R^H .

This gives a significantly shorter proof of Theorem 2.1 in [AS07]. The non-Gaussian stable case also treated in [AS07] cannot be handled this way due to the absence of the decorrelation inequality.

The above technique can be applied in general: assume X can be represented as $X = Y + Z$ with Y and Z being not necessarily independent Gaussian processes. If Z is smoother than Y we can use Theorem 8 to show that Z has a lower small deviation rate and then Li's weak decorrelation inequality [Li99] to show that X and Y have the same small deviations. We summarize this in the following corollary.

Corollary 21. *Let X be a centered Gaussian random variable with values in a normed space $(E, \|\cdot\|)$, where $\|\cdot\| \in \mathbf{N}(\beta, p)$. Let $\kappa > 0$, $\gamma > 0$, and ℓ be a slowly varying function. Assume we can represent X as $X = Y + Z$ with Y and Z also Gaussian random variables in E (not necessarily independent). If*

$$-\log \mathbb{P} [\|Y\| \leq \varepsilon] \sim \kappa \varepsilon^{-\gamma} \ell(\varepsilon)$$

and $Z^{(M)} \in L_2[0, 1]$ for some $M > 1/\gamma - \beta - 1/p$, then

$$-\log \mathbb{P} [\|X\| \leq \varepsilon] \sim \kappa \varepsilon^{-\gamma} \ell(\varepsilon).$$

The same holds for weak asymptotics.

We finish with another example of stable processes.

Example 22. We consider integrated linear fractional stable motions (LFSMs), cf. [ST94]. Let $a, b \in \mathbb{R}$ and $H > 0$ and define

$$X^{a,b}(t) := \int_{\mathbb{R}} a((t-s)_+^{H-1/\alpha} - (-s)_+^{H-1/\alpha}) + b((t-s)_-^{H-1/\tau} - (-s)_-^{H-1/\tau}) dZ(s),$$

where Z is a symmetric α -stable Lévy process, $x_+ := \max(x, 0)$ and $x_- := (-x)_+$. The small ball problem for these processes was treated in [LS05] ($a = 1$, $b = 0$) and in [Aur08] (general a, b) to the end that $X^{a,b}$ has the same small deviation rate (for any $\|\cdot\| \in \mathbf{N}(\beta, p)$) as the corresponding Riemann-Liouville process. From Theorem 6 follows a bound for the small deviation rate of integrated $X^{a,b}$.

References

- [AILvZ08] Frank Aurzada, Ildar A. Ibragimov, Mikhail Lifshits, and Harry van Zanten. Small deviations of smooth stationary Gaussian processes. To appear in: *Theory of Probability and its Applications*, arXiv:0803.4238v1, 2008.
- [AS07] Frank Aurzada and Thomas Simon. Small ball probabilities for stable convolutions. *ESAIM Probab. Stat.*, 11:327–343 (electronic), 2007.

- [Aur08] Frank Aurzada. Small deviations for stable processes via compactness properties of the parameter set. *Statist. Probab. Lett.*, 78(6):577–581, 2008.
- [BGT89] Nick H. Bingham, Charles M. Goldie, and Józef L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [CL03] Xia Chen and Wenbo V. Li. Quadratic functionals and small ball probabilities for the m -fold integrated Brownian motion. *Ann. Probab.*, 31(2):1052–1077, 2003.
- [CS90] Bernd Carl and Irmtraud Stephani. *Entropy, compactness and the approximation of operators*, volume 98 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1990.
- [Der08] Steffen Dereich. The coding complexity of diffusion processes under supremum norm distortion. *Stochastic Process. Appl.*, 118(6):917–937, 2008.
- [DFMS03] Steffen Dereich, Franz Fehrer, Anis Matoussi, and Michael Scheutzow. On the link between small ball probabilities and the quantization problem for Gaussian measures on Banach spaces. *J. Theoret. Probab.*, 16(1):249–265, 2003.
- [DV04] Steffen Dereich and Christian Vormoor. The high resolution vector quantization problem with Orlicz norm distortion. Preprint, 2004.
- [FT04] James Allen Fill and Fred Torcaso. Asymptotic analysis via Mellin transforms for small deviations in L^2 -norm of integrated Brownian sheets. *Probab. Theory Related Fields*, 130(2):259–288, 2004.
- [Gao08] Fuchang Gao. Entropy estimate for k -monotone functions via small ball probability of integrated Brownian motion. *Electron. Commun. Probab.*, 13:121–130, 2008.
- [GHLT03] Fuchang Gao, Jan Hannig, Tzong-Yow Lee, and Fred Torcaso. Laplace transforms via Hadamard factorization. *Electron. J. Probab.*, 8:no. 13, 20 pp. (electronic), 2003.
- [GHT03] Fuchang Gao, Jan Hannig, and Fred Torcaso. Integrated Brownian motions and exact L_2 -small balls. *Ann. Probab.*, 31(3):1320–1337, 2003.
- [GLP03] Siegfried Graf, Harald Luschgy, and Gilles Pagès. Functional quantization and small ball probabilities for Gaussian processes. *J. Theoret. Probab.*, 16(4):1047–1062 (2004), 2003.

- [KL93] James Kuelbs and Wenbo V. Li. Metric entropy and the small ball problem for Gaussian measures. *J. Funct. Anal.*, 116(1):133–157, 1993.
- [KMR06] Stanisław Kwapień, Michael B. Marcus, and Jan Rosiński. Two results on continuity and boundedness of stochastic convolutions. *Ann. Inst. H. Poincaré Probab. Statist.*, 42(5):553–566, 2006.
- [KS98] Davar Khoshnevisan and Zhan Shi. Chung’s law for integrated Brownian motion. *Trans. Amer. Math. Soc.*, 350(10):4253–4264, 1998.
- [Li99] Wenbo V. Li. A Gaussian correlation inequality and its applications to small ball probabilities. *Electron. Comm. Probab.*, 4:111–118 (electronic), 1999.
- [Lif95] Mikhail Lifshits. *Gaussian Random Functions*. Mathematics and Its Applications. Kluwer Academic Publishers, Dordrecht, 1995.
- [Lif08] Mikhail Lifshits. Bibliography on small deviation probabilities. available from: <http://www.proba.jussieu.fr/pageperso/smalldev/biblio.html>, Dec. 2008.
- [LL99] Wenbo V. Li and Werner Linde. Approximation, metric entropy and small ball estimates for Gaussian measures. *Ann. Probab.*, 27(3):1556–1578, 1999.
- [LP08] Harald Luschgy and Gilles Pagès. Functional quantization rate and mean regularity of processes with an application to Lévy processes. *Ann. Appl. Probab.*, 18(2):427–469, 2008.
- [LS01] Wenbo V. Li and Qi-Man Shao. Gaussian processes: inequalities, small ball probabilities and applications. In *Stochastic processes: theory and methods*, volume 19 of *Handbook of Statist.*, pages 533–597, Amsterdam, 2001. North-Holland.
- [LS05] Mikhail A. Lifshits and Thomas Simon. Small deviations for fractional stable processes. *Ann. Inst. H. Poincaré Probab. Statist.*, 41(4):725–752, 2005.
- [Naz03] Alexander I. Nazarov. On the sharp constant in the small ball asymptotics of some Gaussian processes under L_2 -norm. *J. Math. Sci. (N. Y.)*, 117(3):4185–4210, 2003. Nonlinear problems and function theory.
- [NN04a] Alexander I. Nazarov and Yakov Yu. Nikitin. Exact L_2 -small ball behavior of integrated Gaussian processes and spectral asymptotics of boundary value problems. *Probab. Theory Related Fields*, 129(4):469–494, 2004.

- [NN04b] Alexander I. Nazarov and Yakov Yu. Nikitin. Logarithmic asymptotics of small deviations in the L_2 -norm for some fractional Gaussian processes. *Teor. Veroyatn. Primen.*, 49(4):695–711, 2004.
- [Sam98] Gennady Samorodnitsky. Lower tails of self-similar stable processes. *Bernoulli*, 4(1):127–142, 1998.
- [SKM93] Stefan G. Samko, Anatoly A. Kilbas, and Oleg I. Marichev. *Fractional integrals and derivatives*. Gordon and Breach Science Publishers, Yverdon, 1993.
- [ST94] Gennady Samorodnitsky and Murad S. Taqqu. *Stable non-Gaussian random processes*. Stochastic Modeling. Chapman & Hall, New York, 1994.